

SNORING & SLEEP APNEA QUESTIONNAIRE Epworth Sleepiness Scale**

DOB

Date

Gavin Setzen, MD Lawrence S. Kaufman, MD Diplomates American Board of Otolaryngology Fellows American College of Surgeons O = would never

Siobhan Kuhar, MD, PhD

Diplomate American Board of Otolaryngology American Board of Sleep Medicine Fellow

American College of Surgeons American Academy of Otolaryngic Allergy

American Academy of Otolaryngic Allergy

Nora Perkins, MD

Diplomate

American Board of Otolaryngology

John Gavin, MD, FAAP

Fellow American Academy of Pediatrics

Robert Adelson, MD, FACS

Fellow American College of Surgeons

Donna Silvernail, RPA-C Maggie West-Bump, RPA-C Robert Nadratowski, RPA-C Robyn Smith, RPA-C

400 Patroon Creek Boulevard Suite 205

> Albany, New York 12206 Tel: 518 701.2000

Fax: 518 701.2020

www.albanyentandallergy.com

Pediatric & Adult Ear, Nose & Throat Disorders

Sinus, Nasal & Rhinology
Allergy Testing & Treatment
Dizziness & Hearing Loss
Voice, Speech & Swallowing
Snoring & Sleep Apnea
Head & Neck Surgery
Thyroid & Parathyroid Surgery
Facial Plastic Surgery
CAT Scan Imaging
Hearing & Balance Testing

Audiology

Marcia Perretta, AuD Deanna Ross, AuD Tricia Brown, AuD

CAT Scan Imaging

Judith Martin, L.RT, CT.M

Speech Pathology Susan C. Hare, MS, CCC-SLP

Using the scale below, please rank the following activities: 0 = would never doze; 1 = slight chance of dozing; 2 = moderate chance of dozing; 3 = high chance of dozing	
Activity	Chance of Dozing
Sitting and reading	
Watching TV	
Sitting inactive in a public place (e.g. meeting or theater)	
As a passenger in a car for 1 hour without a break	
Lying down in the afternoon when circumstances permit	
Sitting and talking to someone	
Sitting quietly after lunch without alcohol	
In a car, while stopped for a few minutes in traffic	
Total	
*Epworth Sleepiness Scale successfully discriminates popula sleepiness. Individual results may have a wide range and can or determine severity of disease. Obstructive sleep apnea patiseverity of RDI demonstrate:	not be used to exclude
\Box None (mean RDI= 0) an ESS of 8.0 = 3.5	

seventy of RDI demonstrate:	
□ None (mean RDI= 0)	an ESS of $8.0 = 3.5$
□ Mild (mean RDI = 12.1) an ESS of $11.4 = 4.2$	
\square Moderate (mean RDI = 34.8) an ESS of 13.0 = 4.7	
☐ Severe (mean RDI =	56.6) an ESS of 16.2 = 3.3
I have reviewed the above inform	nation with the patient.
Gavin Setzen, MD	Lawrence S. Kaufman, MD
Siobhan Kuhar, MD	Nora Perkins, MD
Robert Adelson, MD	Maggie West-Bump, RPA-C
Maggie West – Bump, RPA-C	Robert Nadratowski, RPA-C
Date:	
Page 1 of 1	Rev: Mar-13